

Your guide to treating
your chronic GVHD
with REZUROCK



REZUROCK[®]

(belumosudil) tablets

When **CHRONIC GVHD** continues to affect your life

ROCK ON with REZUROCK

Bringing balance back to you and your immune system.

GVHD, graft-versus-host disease.

USE

REZUROCK[®] (belumosudil) is a prescription medicine used to treat adults and children 12 years of age and older with chronic graft-versus-host disease (chronic GVHD) after you have received at least 2 prior treatments (systemic therapy) and they did not work. It is not known if REZUROCK is safe and effective in children less than 12 years old.

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney or liver problems.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

Chronic GVHD throws your immune system **OUT OF BALANCE**

Learning about chronic GVHD can help you understand what's happening in your body and why. It can also help you make **IMPORTANT TREATMENT DECISIONS WITH YOUR HEALTH CARE TEAM.**




Chronic GVHD is a common and serious complication that can happen **AFTER A BLOOD STEM CELL TRANSPLANT.**



It occurs **IN ABOUT HALF** of people who receive stem cells from a donor.

CHRONIC GVHD HAPPENS MOST OFTEN IN PEOPLE WHO

- Received a blood stem cell transplant that was mismatched or unrelated
- Received a transplant with donor cells from the bloodstream rather than donor cells collected from bone marrow or umbilical cord blood
- Had acute GVHD
- Are male and received cells from a female
- Are older



During a blood stem cell transplant, **stem cells from a donor (the graft)** are added to **your body (the host)**.

DONOR CELLS MAY ATTACK

**YOUR BODY'S CELLS AND ORGANS
BECAUSE THEY SEE THEM AS A THREAT.**

When this happens, it can lead to chronic GVHD. Chronic GVHD causes your immune system to become **UNBALANCED**, creating too many of some cells and molecules, and not enough of others. This leads to inflammation (swelling) and fibrosis (hardening, scarring), which can damage your organs.

Chronic GVHD is DIFFERENT FOR EVERYONE

Chronic GVHD can happen at any time after your transplant. It can affect your body and how you feel in many ways, so **IT IS IMPORTANT TO RECOGNIZE SYMPTOMS EARLY** and discuss them with your health care team.

CHRONIC GVHD CAN AFFECT MANY DIFFERENT PARTS OF YOUR BODY, SUCH AS



Skin



Eyes



Mouth



Joints and
muscles



Stomach
and
intestines



Liver



Lungs

Common symptoms of chronic GVHD include

- Skin rashes or changes in skin color
- Skin thickening or tightening
- Stiff joints and trouble using your fingers, hands, elbows, knees or ankles
- Dry eyes or vision changes
- Chronic cough, shortness of breath or difficulty breathing
- Nausea, vomiting, diarrhea or loss of appetite
- General decline in health, including fatigue

SOME PEOPLE ALSO HAVE



Lack of energy



Reduced ability to
do daily tasks

Know your options FOR CHRONIC GVHD

Management of chronic GVHD

Management of chronic GVHD is based on which of your organs are affected, the severity of your symptoms and the impact of the disease on your daily life.

Medicines that treat chronic GVHD

Health care providers may try different medicines, including immunosuppressants, to help treat chronic GVHD. However, for some people that might not be enough, and they will need a different approach.



If you have chronic GVHD and are still experiencing symptoms, **TALK WITH YOUR HEALTH CARE TEAM ABOUT YOUR TREATMENT OPTIONS.**



YOU'RE NOT ALONE


Click or Tap to hear other people with chronic GVHD talk about their condition.



REZUROCK WORKS DIFFERENTLY

REZUROCK[®]
(belumosudil) tablets

REZUROCK can help **REBALANCE** your immune system. It is the only approved medicine for chronic GVHD that blocks the ROCK2 pathway in your body. This may reduce the **INFLAMMATION** and **FIBROSIS** that can happen with chronic GVHD.



IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- are pregnant or plan to become pregnant. REZUROCK can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with REZUROCK. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with REZUROCK.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

A different pathway to TREATING CHRONIC GVHD


REZUROCK[®]
(belumosudil) tablets

About the ROCK2 pathway

Many systems in the body work through what are called “**PATHWAYS.**” These pathways are a series of actions that take place among cells and molecules within the body. One such pathway is called “**ROCK2.**”

In people with chronic GVHD, this ROCK2 pathway is too active. This leads to a decrease in cells and molecules that help keep the immune system in check and an increase in cells and molecules that cause inflammation and fibrosis. The result is an immune system that is **OUT OF BALANCE.**



INFLAMMATION is swelling that occurs because your body is trying to protect itself. Inflammation that goes on for a long time, like in chronic GVHD, may cause damage to your body.



FIBROSIS is the growth of scar tissue that hardens and keeps your organs from working properly. In some people, fibrosis happens when inflammation continues for a long time. It is also possible for scar tissue to form without any inflammation.

Go to [REZUROCK.com](https://www.rezurock.com) to watch a video about **HOW REZUROCK WORKS.**

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- **Females** who can become pregnant should use effective birth control during treatment with REZUROCK and for at least 1 week after the last dose.
- **Males** with female partners who can become pregnant should use effective birth control during treatment with REZUROCK and for at least 1 week after the last dose.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

HOW REZUROCK CAN HELP YOU...

REZUROCK[®]
(belumosudil) tablets

REZUROCK is for patients 12 years and older with chronic graft-versus-host disease (chronic GVHD) after 2 prior treatments did not work.

REZUROCK was evaluated in 2 clinical studies that included a total of 186 people with chronic GVHD. One study, called the ROCKstar study, had 132 people. The other study, called the KD025-208 study, had 54 people.

DIFFERENT TYPES OF PEOPLE TOOK REZUROCK IN THE STUDIES, SUCH AS

- People who tried immunosuppressants, such as steroids, and other medicines for chronic GVHD that failed to control their symptoms
- People with mild, moderate and severe chronic GVHD
- People with chronic GVHD in 1 or more organs

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you: (cont)

- are breastfeeding or plan to breastfeed. It is not known if REZUROCK passes into breast milk. Do not breastfeed during treatment with REZUROCK and for at least 1 week after the last dose.

Please see additional Important Safety Information throughout. Please see full [Prescribing Information](#).



ROCK ON

REZUROCK[®]
(belumosudil) tablets

75%

(that's 3 out of every 4 people)

of people in the ROCKstar study saw improvement in their chronic GVHD.

- More than half of people who saw improvement did so between **WEEKS 4 AND 8**
- Almost everyone who saw improvement did so by **WEEK 24**
- Medicine(s) may affect everyone differently. Some people may take longer to respond, especially if they have severe disease caused by fibrosis

RESPONSE WAS SEEN IN VARIOUS ORGANS



Skin



Eyes



Mouth



Joints and muscles



GI tract



Liver



Lungs



Esophagus

Some people taking REZUROCK had

- **A REDUCED USE OF STEROIDS** and other immunosuppressants
- **QUALITY-OF-LIFE IMPROVEMENTS**, reported in an exploratory analysis^a

GI, gastrointestinal.

^aAn exploratory analysis provides a better understanding of a problem but is not proof. It is difficult to measure quality of life because every person is different and will have a unique experience with treatment.

IMPORTANT SAFETY INFORMATION

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REZUROCK may affect the way other medicines work, and other medicines may affect the way REZUROCK works.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

WHAT SIDE EFFECTS should I be aware of with REZUROCK?

REZUROCK[®]
(belumosudil) tablets



The **most common side effects** reported by people taking REZUROCK in 2 clinical studies were

- Infections
- Tiredness or weakness
- Nausea
- Diarrhea
- Shortness of breath
- Cough
- Swelling
- Bleeding
- Stomach (abdominal) pain
- Muscle or bone pain
- Headache
- High blood pressure

REZUROCK may affect fertility in males and females. Talk with your health care provider if this is a concern for you.

These are not all the possible side effects of REZUROCK.

IMPORTANT SAFETY INFORMATION

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

Additional INFORMATION

REZUROCK[®]
(belumosudil) tablets



In clinical studies, there were **NO NEW CASES** of cytomegalovirus (CMV) infection and only 1 case of recurring CMV infection.

CMV is a common serious infection that can happen in people who have had a blood stem cell transplant.



About **4% OF PEOPLE** (that's 1 out of every 25 people) in clinical studies had serious cytopenia (site-oh-pee-nia), a condition that causes a drop in levels of certain blood cells.

Talk with your health care team to find out if REZUROCK is right for you.



If you have chronic GVHD, talk with your health care provider about staying on REZUROCK as long as needed. **THIS IS VERY IMPORTANT.**

Be sure to tell your health care provider about any side effects you may have. They may be able to help you find ways to manage them.

Please see Important Safety Information throughout.
Please see full [Prescribing Information](#).

How to TAKE REZUROCK

REZUROCK[®]
(belumosudil) tablets

The REZUROCK 200-mg tablet is taken by mouth **once a day with a meal.**



The REZUROCK tablet should be swallowed whole with a glass of water.
DO NOT CUT, CRUSH OR CHEW THE TABLET.



Take REZUROCK at approximately the same time each day with a meal.
A MEAL IS NOT A SNACK. It is about the same amount of calories as you might have for lunch or dinner.



If you miss a dose of REZUROCK, take it as soon as you remember on the same day. Take your next dose of REZUROCK at your regular time on the next day.
DO NOT TAKE EXTRA DOSES OF REZUROCK to make up for a missed dose.

Take REZUROCK as prescribed by your health care team

- Always talk with your health care team about any changes to your medications
- Do not start or stop taking any medicines without discussing it first with your health care team. If you experience a side effect, be sure to tell your health care team right away
- Always let your health care team know about all the medications you are taking

IMPORTANT SAFETY INFORMATION

How should I take REZUROCK?

- Take REZUROCK exactly as your healthcare provider tells you to take it.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

REZUROCK[®]

(belumosudil) tablets

How to store REZUROCK

Store REZUROCK at room temperature, ranging from 68 °F to 77 °F (20 °C–25 °C). REZUROCK should be stored in its original container to protect it from moisture. Make sure to close the cap tightly each time after taking your tablet. Do not discard the drying packet inside the bottle.

If you are taking REZUROCK, make sure to tell your health care team about all of the prescription and over-the-counter medicines (including proton pump inhibitors, or PPIs), vitamins and herbal products you are taking because they may affect how REZUROCK works.

IMPORTANT SAFETY INFORMATION

How should I take REZUROCK? (cont)

- Do not change your dose or stop taking REZUROCK without first talking to your healthcare provider.



THERE'S MORE TO LEARN

Click or Tap for REZUROCK updates or visit REZUROCK.com.



Please see additional Important Safety Information throughout.
Please see full [Prescribing Information](#).

How to GET REZUROCK


REZUROCK[®]
(belumosudil) tablets

Through Kadmon ASSIST, we strive to help and support people with chronic GVHD, and their caregivers, throughout their treatment journey.



Visit [REZUROCK.com](https://www.rezurock.com) or call **1-844-KADMON1 (523-6661)** to find out if you're eligible to receive help getting started on REZUROCK treatment.

Talk with your health care team about enrolling in Kadmon ASSIST.

Kadmon ASSIST is available to help eligible people

- ✓ Determine insurance coverage and costs for REZUROCK
- ✓ Find a savings program they may qualify for
- ✓ Have medicine delivered directly by a specialty pharmacy
- ✓ Get support from an adherence nurse about treatment with REZUROCK

Kadmon does not provide medical advice, diagnosis or treatment recommendations.

SUPPORT STARTS HERE

REZUROCK[®]
(belumosudil) tablets

Learn more about the Kadmon ASSIST support programs designed for people who need treatment for chronic GVHD at [REZUROCK.com](https://www.rezurock.com), or call **1-844-KADMON1 (523-6661)**.

Our Kadmon ASSIST team is available to help you **Monday through Friday, 8 AM-8 PM ET**.

Eligible commercially or privately insured people can receive REZUROCK for

**AS LOW AS
\$0 PER MONTH**

through the Kadmon ASSIST Commercial Co-pay Savings Program.^a



^aPatient Terms and Conditions: The Kadmon ASSIST Commercial Co-pay Savings Program provides co-pay/coinsurance support for out-of-pocket costs on REZUROCK[®] (belumosudil) tablets prescriptions. A yearly maximum benefit applies. Limit one 30-day supply per 30 days. This program is not health insurance. This program is for commercially or privately insured patients only; uninsured or cash-paying patients are not eligible. Patients are not eligible if prescriptions are paid, in whole or in part, by any state- or federally funded programs, including, but not limited to, Medicare (including Part D, even in the coverage gap) or Medicaid, Medigap, VA, DOD, TriCare, private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs, or where prohibited by law. The co-pay program may not be combined with any other rebate, coupon or offer. Kadmon Pharmaceuticals, LLC, reserves the right to rescind, revoke or amend this offer at any time without further notice. Any savings provided by the co-pay program may vary depending on patients' out-of-pocket costs. Card is valid through December 31 of the year of activation. On January 1 of the following year, the card automatically resets and is subject to annual limits if the prescription benefit remains the same. Upon registration, patients receive all program details.

USE AND IMPORTANT SAFETY INFORMATION



USE

REZUROCK® (belumosudil) is a prescription medicine used to treat adults and children 12 years of age and older with chronic graft-versus-host disease (chronic GVHD) after you have received at least 2 prior treatments (systemic therapy) and they did not work. It is not known if REZUROCK is safe and effective in children less than 12 years old.

IMPORTANT SAFETY INFORMATION

Before taking REZUROCK, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney or liver problems.
- are pregnant or plan to become pregnant. REZUROCK can harm your unborn baby. If you are able to become pregnant, your healthcare provider will do a pregnancy test before starting treatment with REZUROCK. Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with REZUROCK.
 - **Females** who can become pregnant should use effective birth control during treatment with REZUROCK and for at least 1 week after the last dose.
 - **Males** with female partners who can become pregnant should use effective birth control during treatment with REZUROCK and for at least 1 week after the last dose.
- are breastfeeding or plan to breastfeed. It is not known if REZUROCK passes into breast milk. Do not breastfeed during treatment with REZUROCK and for at least 1 week after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REZUROCK may affect the way other medicines work, and other medicines may affect the way REZUROCK works.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take REZUROCK?

- Take REZUROCK exactly as your healthcare provider tells you to take it.
- Do not change your dose or stop taking REZUROCK without first talking to your healthcare provider.
- Take REZUROCK 1 time a day with a meal.
- Take REZUROCK at about the same time each day.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

IMPORTANT SAFETY INFORMATION (cont)

How should I take REZUROCK? (cont)

- Swallow REZUROCK tablets whole with a glass of water.
- Do not cut, crush, or chew REZUROCK tablets.
- Your healthcare provider will do blood tests to check your liver at least 1 time a month during treatment with REZUROCK.
- If you miss a dose of REZUROCK, take it as soon as you remember on the same day. Take your next dose of REZUROCK at your regular time on the next day. Do not take extra doses of REZUROCK to make up for a missed dose.
- If you take too much REZUROCK, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of REZUROCK?

The most common side effects of REZUROCK include:

- infections
- swelling
- tiredness or weakness
- bleeding
- nausea
- stomach (abdominal) pain
- diarrhea
- muscle or bone pain
- shortness of breath
- headache
- cough
- high blood pressure

Your healthcare provider may change your dose of REZUROCK, temporarily stop, or permanently stop treatment with REZUROCK if you have certain side effects.

REZUROCK may affect fertility in males and females. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of REZUROCK. Call your doctor for medical advice about side effects.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch or call **1-800-FDA-1088**. You may also contact Kadmon Pharmaceuticals, LLC, a Sanofi Company, at **1-800-633-1610** to report side effects.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

ROCK ON WITH REZUROCK



- REZUROCK is the only approved medicine that blocks the ROCK2 pathway to help **REBALANCE** your immune system. It **MAY REDUCE THE INFLAMMATION AND FIBROSIS** that can happen with chronic GVHD
- **75% OF PEOPLE** (that's 3 out of every 4 people) in the ROCKstar study saw **IMPROVEMENT IN THEIR CHRONIC GVHD**
- More than half of people who saw improvement did so between **WEEKS 4 AND 8**
- Some people had a **REDUCED USE OF STEROIDS AND OTHER IMMUNOSUPPRESSANTS**
- **QUALITY-OF-LIFE IMPROVEMENTS SEEN BY SOME PEOPLE**, reported in an exploratory analysis^a
- REZUROCK is a **TABLET**, taken by mouth, that is approved for **ONCE-DAILY DOSING**



Through the Kadmon ASSIST program, we strive to help and support people with chronic GVHD, and their caregivers, throughout their treatment journey.

Ready to ROCK ON? Ask your health care team if REZUROCK is right for you. For more information about REZUROCK, chronic GVHD and Kadmon ASSIST, visit [REZUROCK.com](https://www.rezurock.com).

^aAn exploratory analysis provides a better understanding of a problem but is not proof. It is difficult to measure quality of life because every person is different and will have a unique experience with treatment.

USE

REZUROCK[®] (belumosudil) is a prescription medicine used to treat adults and children 12 years of age and older with chronic graft-versus-host disease (chronic GVHD) after you have received at least 2 prior treatments (systemic therapy) and they did not work. It is not known if REZUROCK is safe and effective in children less than 12 years old.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of REZUROCK?

REZUROCK may affect fertility in males and females. Talk to your healthcare provider if this is a concern for you.

Please see additional Important Safety Information throughout.

Please see full [Prescribing Information](#).

Sanofi does not provide medical advice, diagnosis or treatment. The health information contained herein is provided for general educational purposes only. Your health care professional is the best source of information regarding your health. Please consult your health care professional if you have any questions about your health or treatment.



© 2023 Kadmon Pharmaceuticals, LLC. All Rights Reserved.
REZUROCK, Kadmon and Sanofi are registered trademarks of Sanofi or an affiliate.
Kadmon ASSIST is owned by Sanofi or an affiliate.
MAT-US-2203537-v3.0-11/2023

REZUROCK[®]
(belumosudil) tablets